



# Primordial pseudobulges in isolated galaxies?

Mirian Fernández Lorenzo  
IAA-CSIC

Collaborators: J. Sulentic, L. Verdes-Montenegro, Blasco-Herrera, J., M. Argudo-Fernández, Ramirez-Moreta, P., Garrido, J., J.E. Ruiz, S. Sánchez-Expósito, Santánder-Vela, J.D

# Central parts of galaxies

**Classical bulge** – built through rapid/violent processes (e.g. major mergers)

- ★ Old stellar populations
- ★ Dynamically supported by velocity dispersion
- ★ Follows the same relations than E/S0

*Surface brightness profiles: ~ De Vaucouleurs,  $n > 2$*

**Pseudobulge** – built through slow/secular processes (e.g. gas infall, star formation; Kormendy & Kennicutt 2004)

- ★ Young stellar populations and SF
- ★ Rotation motions
- ★ Disky structures – memory of their disk origin
- ★ Nuclear bars and rings

*Surface brightness profiles: ~ Exponential,  $n < 2$*

***Isolated galaxies – Bulge evolution mainly driven by internal processes***

# The AMIGA project

## Analysis of the interstellar Medium of Isolated GALaxies

Started in 2003 @IAA (PI: L. Verdes-Montenegro)

- Catalogue of Isolated Galaxies (CIG) - 1051 (Karachentseva 1973)
  - Very restrictive selection criteria - No major **tidal interaction** within the last ~3 Gyr
  - Better than **field** (pairs, triplets)
- Goal: to quantify the properties of the CIG interstellar medium
  - UV to radiocontinuum study - different properties at all wavelength
  - To separate effects from “**Nature**” and “**Nurture**”
- Main results in the optical:
  - A higher fraction of spirals – only 15% early-types
  - Most Sb-Sc spirals present pseudobulges

# Central parts of galaxies

AMIGA project: Analysis of the interstellar Medium of Isolated Galaxies


Catalogue of Isolated Galaxies (CIG) - 1051 (Karachentseva 1973)

No major **tidal interaction** within the last  $\sim 3$  Gyr

Main results in the optical:

- A higher fraction of spirals – only 15% early-types
- Most Sb-Sc spirals present pseudobulges

Durbala+ (2008)



# Central parts of isolated galaxies

## Bulge classification

(Fernández Lorenzo+ 2014)

Increase the sample (94) in Durbala+ (2008) to all AMIGA spirals in SDSS

## Sample selection:

★ Galaxies that follow the isolation criteria of Argudo-Fernández+ (2013)

★ Completeness criteria:  $\text{mag B} < 15.3$  ( $\sim \text{mag r} < 14.5$ )

⇒ Final sample: 298 galaxies



★ Bulge/disk/bar decomposition with GALFIT in the i-band

# Central parts of isolated galaxies

## Bulge classification

(Fernández Lorenzo+ 2014)

Final sample: 189 galaxies (residuals in the center lower than 10%)





# Central parts of isolated galaxies

## Bulge colors

(Fernández Lorenzo+ 2014)

(g-i) colors as indicative of the **stellar populations**

- ★ Disk fits in g and i-bands: independent of fixed parameters
- ★ Bulge colors from galfit more than  $3\sigma$  redder than the red sequence: a change in the disk inside the bulge because the bulge formation and evolution?

(g-i) bulge colors: **aperture photometry** with ellipse

- ★ Galaxies fitted in the i-band
- ★ Aperture magnitudes in r and g-bands: ellipticity and position angle of isophotal aperture equal to the i-band values.

# Central parts of isolated galaxies


## Bulge colors

(Fernández Lorenzo+ 2014)

Color-magnitude relation of galaxies in the Nair & Abraham (2010) sample

**63%** of bulge in the red sequence

**58%** of high-BT and **66%** of low-BT pseudobulges in the red sequence






# Central parts of isolated galaxies

## Bulge colors

(Fernández Lorenzo+ 2014)

- ★ Corrected by Galactic extinction, k-correction and reddening (inclination)
- ★ What about a **full reddening** correction?
- ★ We used  $A_v$  from starlight (Cid-Fernandes et al. 2005) for galaxies with spectra



# Central parts of isolated galaxies

## Bulge colors

(Fernández Lorenzo+ 2014)

Starburst99 (Leitherer et al. 1999)

Mean bulge stellar mass =  $3 \times 10^9 M_{\odot}$

Two simulations (IMF of Kroupa,  $Z=0.008$ ):

1) **Instantaneous burst** of  $1.5 \times 10^9 M_{\odot}$


2) **Continuous SF** of  $0.2 M_{\odot} \text{ yr}^{-1}$

After 8 Gyr:

★ Instantaneous burst:  $(g-i) = 1.13$

★ Continuous SF:  $(g-i) = 0.49$

★ Combined model:  $(g-i) = 0.59$




# Central parts of isolated galaxies

Dependence with the environment

(Fernández Lorenzo+ 2014)

(g-i) pseudobulge color Vs  $Q_{kar}$



- ★ **Red pseudobulges** distributed in all range
- ★ **Blue pseudobulges** tend to be located at higher values of  $Q_{kar}$



# Central parts of isolated galaxies

## Galaxies without bar

- ⇒ 32% of High-BT pseudobulges
- ⇒ Most classical bulges
- ⇒ 62% of Low-BT pseudobulges





# Central parts of isolated galaxies

## Galaxies with bar

⇒ 68% of High-BT pseudobulges

⇒ 38% of Low-BT pseudobulges






# Central parts of isolated galaxies

## Galaxies without bar

- ⇒ 32% of High-BT pseudobulges
- ⇒ Most classical bulges
- ⇒ 62% of Low-BT pseudobulges






# Central parts of isolated galaxies

## Galaxies with bar

⇒ 68% of High-BT pseudobulges

⇒ 38% of Low-BT pseudobulges



**Thanks!**